

5/23/16 "Live as if you were to die tomorrow. Learn as if you were to live forever."
-Mahatma Gandhi

HW: Study for the Final
Test 3 Thursday 6/2

AIM: What is on the Final?

Warm Up:

1)

$$1) \text{ Length} = \int_a^b \sqrt{1 + (f'(x))^2} \, dx$$

$$f(x) = x^2$$

$$f'(x) = 2x$$

$$\text{Length} = \int_0^2 \sqrt{1 + (2x)^2} \, dx = 4.647$$

$$g(x) = x^3$$

$$g'(x) = 3x^2$$

$$\text{Length} = \int_0^2 \sqrt{1 + (3x^2)^2} \, dx = 8.630$$

$y = x^3$ is longer over $[0, 2]$

2) Antiderivative:

a) $\int 3x^4 + 2 \, dx$

$$f(x) = \frac{3x^5}{5} + 2x + c$$

b) $\int (5\sqrt{x} - 3x) \, dx$

$$\int (5x^{\frac{1}{2}} - 3x) \, dx$$


$$f(x) = \frac{5x^{\frac{3}{2}}}{\frac{3}{2}} - \frac{3x^2}{2} + c$$

$$= \frac{10}{3}x^{\frac{3}{2}} - \frac{3}{2}x^2 + c$$

$$= \frac{10}{3}\sqrt{x^3} - \frac{3}{2}x^2 + c$$

$$\begin{aligned} 3) \quad a) \quad & \int_0^{\frac{\pi}{2}} \cos(x) \, dx \\ &= \sin x \Big|_0^{\frac{\pi}{2}} \\ &= \sin\left(\frac{\pi}{2}\right) - \sin(0) \\ &= 1 - 0 \\ &= \boxed{1} \end{aligned}$$

$$\begin{aligned} b) \quad & \int_1^2 (3x + x^3) \, dx \\ &= \left[\frac{3x^2}{2} + \frac{x^4}{4} \right]_1^2 \\ &= \left(\frac{3(2)^2}{2} + \frac{2^4}{4} \right) - \left(\frac{3(1)^2}{2} + \frac{1^4}{4} \right) \\ &= (6 + 4) - \left(\frac{3}{2} + \frac{1}{4} \right) \\ &= 10 - \frac{7}{4} \\ &= \boxed{\frac{33}{4}} \end{aligned}$$



NORMAL FLOAT AUTO REAL DEGREE MP

$$\int_0^2 (-x^2 + 4 - ((x-2)^2)) dx$$

2.666666667

Ans►Frac

 $\frac{8}{3}$

5) $y = x^2 - 4$ $y = 0$ (x-axis)

$$V = \pi \int_a^b f(x)^2 dx$$

NORMAL FLOAT AUTO REAL DEGREE MP

$$\int_{-2}^2 ((x^2 - 4)^2) dx$$

34.13333333

Ans►Frac

 $\frac{512}{15}$

$$V = \frac{512}{15} \uparrow \text{units}^3$$

6) $y = \sqrt{x-2}$ $y = 0$ $x = 5$

NORMAL FLOAT AUTO REAL DEGREE MP

$$\int_2^5 ((\sqrt{x-2})^2) dx$$

4.5

$$4.5 \uparrow \text{units}^3$$

8) $y^2 - 2x = 3$ what is $\frac{dy}{dx}$ @ $(3,3)$

$\frac{d}{dx}(y^2 - 2x = 3)$ ← Implicit derivative Evaluate derivative

$$2y \frac{dy}{dx} - 2 = 0$$

$$\frac{2y \frac{dy}{dx}}{2y} = \frac{2}{2y} \rightarrow \frac{dy}{dx} = \frac{2}{2y} = \frac{1}{y}$$

Evaluate @ $(3,3)$

$$\frac{dy}{dx} \Big|_{(3,3)} = \left(\frac{1}{3} \right)$$

$$9) a) \lim_{x \rightarrow 0} \frac{24x^3 - 3x}{12x^3 + x} = \frac{24(0)^3 - 3(0)}{12(0)^3 + (0)} = \frac{0}{0} \text{ Ind.}$$

Try to reduce

$$= \frac{\cancel{3}x(8x^2 - 1)}{\cancel{x}(12x^2 + 1)} = \frac{24x^2 - 3}{12x^2 + 1} \quad \text{Try to plug in again}$$

$$= \frac{24(0)^2 - 3}{12(0)^2 + 1} = \frac{-3}{1}$$

$$\lim_{x \rightarrow 0} \text{ is } (-3)$$

$$b) \lim_{x \rightarrow \infty} \frac{24x^3 - 3x}{12x^3 + x}$$

degrees	limit
$N > D$	∞ or $-\infty$
$D > N$	0
$N = D$	coefficients

Look at
degrees (highest
exponents)

$$\frac{24}{12} = \boxed{2}$$

10) $y = \frac{x+2}{2x-1}$

Quotient Rule

tangent when
 $x = 3$

need slope

$$y' = \frac{(2x-1)(1) - (x+2)(2)}{(2x-1)^2}$$

$$y' @ x=3 = \frac{(2(3)-1)(1) - (3+2)(2)}{(2(3)-1)^2} = \frac{-5}{25}$$

Point:

$$x=3 \quad y = \frac{3+2}{2(3)-1} = \frac{5}{5} = 1$$

Slope of tangent = $\frac{-1}{5}$

$(3, 1)$

Equation of tangent line


$$y - 1 = -\frac{1}{5}(x - 3)$$

@
 $x=3$

⊗ Normal line is exactly
the same with negative reciprocal
slope.

11) Concave down, the 2nd derivative is negative.

$$y = 6x^2 + \frac{x}{2} + 3 + \frac{6}{x}$$



$$12) \quad f(x) = \frac{x^2-1}{2x} \quad f \quad g$$

$$f'(x) = \frac{2x(2x) - (x^2-1)(2)}{(2x)^2}$$

$$= \frac{4x^2 - 2x^2 + 2}{4x^2} = \frac{2x^2 + 2}{4x^2}$$

Product Rule:
 $f'g + fg'$

Quotient Rule:
 $\frac{gf' - fg'}{(g)^2}$

$$\boxed{\frac{x^2 + 1}{2x^2}}$$

13) Position $x(t) = \frac{1}{2} \sin(t) + \cos(2t)$

Velocity $x'(t) = \frac{1}{2} \cos(t) - 2 \sin(2t)$

Acceleration $x''(t) = -\frac{1}{2} \sin(t) - 4 \cos(2t)$

$$x''\left(\frac{\pi}{2}\right) = -\frac{1}{2} \sin\left(\frac{\pi}{2}\right) - 4 \cos\left(2 \cdot \frac{\pi}{2}\right)$$

$$= -\frac{1}{2}(1) - 4(-1)$$

$$= \boxed{3.5}$$

$$14) f(x) = e^x \ln x$$

$$f'(x) = e^x \cdot \frac{1}{x} + e^x \ln x$$

$$f'(e) = e^e \cdot \frac{1}{e} + e^e \ln e$$

$$= \frac{e^e}{e} + e^e(1) \approx \boxed{20.729}$$

Recall

$$f(x) = e^x \quad f'(x) = e^x$$

$$f(x) = \ln x \quad f'(x) = \frac{1}{x}$$

$$\ln_e e = 1$$


$$\text{Area} = \int_a^b \text{Top} - \text{bottom} \, dx$$

15. The graph of f is shown. Evaluate each integral by interpreting in terms of areas.

- a. $\int_0^2 f(x) dx$ Area between $f(x)$ and x -axis between 0 and 2
 b. $\int_2^5 f(x) dx$
 c. $\int_0^5 f(x) dx$

$$b) = 6$$

$$c) 4 + 6 = 10$$



7) Derivative = $6x^2 - 2x + 3$
(Slope)

Looking for
Antiderivative

$$y' = 6x^2 - 2x + 3$$

$$y = \int 6x^2 - 2x + 3 \, dx$$

$$y = \frac{6x^3}{3} - \frac{2x^2}{2} + 3x + C$$

$$y = 2x^3 - x^2 + 3x + C$$

Passes
through
(1, 3)

$$3 = 2(1)^3 - 1^2 + 3(1) + C$$


$$3 = 2 - 1 + 3 + C$$

$$3 = 4 + C$$

$$\begin{array}{r} -4 \quad -4 \\ \hline \end{array}$$

$$-1 = C$$

$$y = 2x^3 - x^2 + 3x - 1$$

Know

$$y = 13 \quad \frac{dy}{dt} = 0$$

$$z = 12 \quad \frac{dz}{dt} = 3$$

$$x = 5$$

want:

$$\frac{dx}{dt}$$

Find x:

$$x^2 + 12^2 = 13^2$$

$$x = 5$$

$$x^2 + z^2 = y^2$$

$$2x \frac{dx}{dt} + 2z \frac{dz}{dt} = 2y \frac{dy}{dt}$$

$$2(5) \frac{dx}{dt} + 2(12)(3) = 2(13)(0)$$

$$10 \frac{dx}{dt} + 72 = 0$$

$$10 \frac{dx}{dt} = -72$$

$$\frac{dx}{dt} = \frac{-72}{10} = -7.2 \text{ ft/sec}$$

Top of ladder
is sliding down
the wall
@ 7.2 ft/sec

$$18) \frac{2(x+h)^2 - 8(x+h) + 5 - (2x^2 - 8x - 5)}{h}$$

$$\frac{\cancel{2x^2} + 4xh + \cancel{2h^2} - \cancel{8x} - \cancel{8h} + 5 - \cancel{2x^2} + \cancel{8x} + \cancel{5}}{h}$$


$$\frac{4xh + 2h^2 - 8h}{h} = \frac{\cancel{h}(4x + 2h - 8)}{\cancel{h}} = 4x + 2h - 8$$

$h \rightarrow 0$
 $4x - 8$

20. If f and g are the functions shown below. Let $h(x) = f(g(x))$ and $s(x) = f(x)g(x)$.

Find: $h'(1)$ and $s'(1)$

$$g(1) = 1$$



Chain

Product

$$h'(x) = f'(g(x)) \cdot g'(x)$$

$$h'(1) = f'(g(1)) \cdot g'(1)$$

$$= f'(1) \cdot g'(1)$$

$$h'(1) = 2 \cdot (-1) = -2$$

$$s'(x) = f'(x) \cdot g'(x) + f'(x)g(x)$$

$$s'(1) = f'(1) \cdot g'(1) + f'(1) \cdot g(1)$$

$$= (2) \cdot (-1) + (2)(1)$$

$$= -2 + 2$$

$$= 0$$

21. The following table records the values of f , f' , g , and g' at $x=1$, $x=2$, and $x=3$.

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	2	1	2	3
2	5	4	3	4
3	0	6	-1	-2

quotient.

chain

If $n(x) = \frac{f(x)}{g(x)}$, $h(x) = f(g(x))$, find the value of each of the following: a) $n'(2)$ b) $h'(1)$

$$a) n'(x) = \frac{g(x)f'(x) - f(x)g'(x)}{(g(x))^2}$$

$$n'(2) = \frac{g(2) \cdot f'(2) - f(2) \cdot g'(2)}{(g(2))^2} = \frac{(3)(4) - (5)(4)}{3^2} = \left(-\frac{8}{9}\right)$$

$$b) h'(x) = f'(g(x)) \cdot g'(x)$$

$$h'(1) = f'(g(1)) \cdot g'(1)$$

$$= f'(2) \cdot (3)$$

$$= (4)(3)$$

$$h'(1) = 12$$

approaching 1 from left

22. $\lim_{x \rightarrow 1^-} f(x) = -\infty$

23. $\lim_{x \rightarrow 1^+} f(x) = \infty$


24. $\lim_{x \rightarrow 1} f(x) = \text{DNE}$

25. $\lim_{x \rightarrow -3} f(x) = 1$

26. $\lim_{x \rightarrow 3} f(x) = -1$

27. $\lim_{x \rightarrow -\infty} f(x) = \infty$

28. $\lim_{x \rightarrow \infty} f(x) = -2$



$$29) \quad y = 3x^2 - 2x + 1$$

Tangent @
 $x = -1$

Point:

$$y = 3(-1)^2 - 2(-1) + 1$$

$$y = 6$$

$$(-1, 6)$$

Slope: (Derivative)

$$y' = 6x - 2$$

$$y' = 6(-1) - 2$$

$$y' = -8$$

$$y - 6 = -8(x + 1)$$

29) b) Equation of normal line @ $x = -1$

$$y - 6 = \frac{1}{8}(x + 1)$$

Negative
reciprocal slope
of tangent.

$$\begin{aligned} 30) \quad s(t) &= 2t^3 - 15t^2 + 24t - 10 \\ v(t) &= 6t^2 - 30t + 24 \\ a(t) &= 12t - 30 \end{aligned}$$

$$a) \quad s(3) = 2(3)^3 - 15(3)^2 + 24(3) - 10 = \boxed{-19}$$

$$b) \quad v(3) = 6(3)^2 - 30(3) + 24 = \boxed{-12}$$

$$c) \quad a(3) = 12(3) - 30 = \boxed{6}$$

d) Slowing down
b/c $v(3)$ and $a(3)$
have different signs.


e) At rest when $v(t) = 0$

$$\begin{aligned} 0 &= 6t^2 - 30t + 24 \\ &= 6(t^2 - 5t + 4) \\ &= 6(t-4)(t-1) \end{aligned}$$

$t=4 \quad | \quad t=1$

$$\begin{aligned} t &= 4 \\ t &= 1 \end{aligned}$$

f) Moving \rightarrow when $v(t)$ is positive



Moving Right $[0, 1) \cup (4, \infty)$

g) Used to be tough b/c we had to calculate distance over intervals of left and right motion and then add them.

$$\text{Distance} = \int_a^b |v(t)| dt$$

absolute value to get rid of + and - velocity


$$\text{Distance} = \int_0^3 |6t^2 - 30t + 24| dt$$

NORMAL FLOAT AUTO REAL DEGREE HP

$$\int_0^3 (16x^2 - 30x + 24) dx$$

.....31.0000067.....

$$\text{Dist} = 31$$



$$8^2 + 15^2 = z^2$$

$$64 + 225 = z^2$$

$$289 = z^2$$

$$17 = z$$

Have:

$$x=8 \quad \frac{dx}{dt}=4$$

$$y=15 \quad \frac{dy}{dt}=7.5$$

$$z=17 \quad \frac{dz}{dt}=7.5$$

8.5 mph

$$x^2 + y^2 = z^2$$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 2z \frac{dz}{dt}$$

$$2(8)(4) + 2(15)(7.5) = 2(17) \frac{dz}{dt}$$

$$64 + 225 = 34 \frac{dz}{dt}$$

$$\frac{289}{34} = 34 \frac{dz}{dt}$$

8.5 = $\frac{dz}{dt}$

$$32) \quad x^2 - xy = y^2 + 1$$

$$y = \textcircled{1} \quad \underline{QI}$$

$$\frac{d}{dx} \overset{\text{Slope}}{(x^2 - xy = y^2 + 1)}$$

$$2x - \left(x \frac{dy}{dx} + 1y\right) = 2y \frac{dy}{dx}$$

$$2x - x \frac{dy}{dx} - y = 2y \frac{dy}{dx}$$

$$\textcircled{2(2)} - 2 \frac{dy}{dx} \textcircled{-1} = 2(1) \frac{dy}{dx}$$

$$\textcircled{3} - 2 \frac{dy}{dx} = 2 \frac{dy}{dx}$$

$$+ 2 \frac{dy}{dx} \quad + 2 \frac{dy}{dx}$$

$$\frac{3}{4} = \frac{4 \frac{dy}{dx}}{4}$$

$$\frac{dy}{dx} = \frac{3}{4}$$

$$y - 1 = \frac{3}{4}(x - 2)$$

point:

$$x^2 - x(1) = 1^2 + 1$$

$$x^2 - x = 2$$

$$x^2 - x - 2 = 0$$

$$(x-2)(x+1) = 0$$

$$\textcircled{x=2} \quad x=-1$$

QI reject not QI

$$(2, 1)$$

Point

